

119. Umsetzung von Cyclopentadienylmetallcarbonylen mit Thionylhalogeniden – neue Synthese halogenverbrückter Cp*Co-Komplexe, C. Stoll, I.-P. Lorenz, K. Polborn und E.F. Paulus, *Z. Naturforsch.* **1999**, *54b*, 583–590.
120. Halogenation and oxygenation of the metal and oxidation of phosphorus in complexes of the type $M(dppe)_2(N_2)_2$ ($M = Mo, W$) by reaction with thionylhalides, C. Stoll, I.-P. Lorenz, H. Nöth and W. Ponikwar, *J. Organomet. Chem.* **2000**, *602*, 24–28.
121. Chromophores encapsulated in Gold complexes: dpp dyes with novel properties, H. Langhals, M. Limmert, I.-P. Lorenz, P. Mayer, H. Piotrowski and K. Polborn, *Eur. J. Inorg. Chem.* **2000**, 2345–2349.
122. Halogen-Bridged cationic complexes of the type $[\{CpM(CO)_3\}_2X]SbF_6$ ($M = Mo, W$; $X = Cl, I$) and $[\{CpW(CO)_2Cl_2\}_2Cl]SbCl_6$, and the trichloride $CpW(CO)_2Cl_3$, I.-P. Lorenz, R. Schneider und H. Nöth, *Z. Anorg. Allg. Chem.* **2000**, *626*, 1958–1962.
123. Koordination und Reaktivität von Sulfinylimiden $RNSO$ in Cyclopentadienyldicarbonylmangan-Komplexen, I.-P. Lorenz, M. Limmert, K. Polborn and H. Nöth, *Z. Anorg. Allg. Chem.* **2000**, 2313–2317.
124. BF_3 -Stabilized metalated sulfinylimides and sulfuriimides of the type $Cp(NO)_2CrNSE.BF_3$ ($E = O, NSiMe_3$), M. Limmert, I.-P. Lorenz, J. Neubauer, H. Nöth and T. Habereder, *Eur. J. Inorg. Chem.* **2001**, 1593–1597.
125. Organometalated halonium salts of the type $[\{Cp(CO)_2Fe\}_2X]SbY_6$ ($X = Cl, Br, I; Y = F, Cl$), R. Schneider, I.-P. Lorenz, H. Nöth and W. Ponikwar, *Z. Anorg. Allg. Chem.* **2001**, *627*, 1775–1781.
126. Synthesis and structure of di- and triferristibonium salts of the type $[Fp_2SbPh_2]_2FeX_4$ and $[Fp_3SbR]_2FeX_4$ ($Fp = CpFe(CO)_2$, $R = Ph, Cl$; $X = Cl, Br, I$), I.-P. Lorenz, R. Schneider, H. Nöth, K. Polborn and J. Breunig, *Z. Naturforsch.* **2001**, *56b*, 671–679.
127. New organometalated phosphinato-O,O \square ligands in the planar eight-membered ring $[\{Cp(CO)_2\}'BuPO_2FeCl_2]_2$, I.-P. Lorenz, M. Maier and K. Polborn, *Eur. J. Inorg. Chem.* **2002**, 327–330.
128. DPP dyes as ligands in copper, silver, and gold complexes, I.-P. Lorenz, M. Limmert, P. Mayer, H. Piotrowski, H. Langhals, M. Poppe and K. Polborn, *Chem. Eur. J.* **2002**, *8*, 4047–4055.
129. Synthesis of the dimeric μ_2 -O aminoalkoxy complexes of rhenium $[(CO)_3Re(\mu_2$ -O $\cap NH_2)]_2$ and their condensation products with acetone $[(CO)_3Re(\mu_2$ -O $\cap NH=CMe_2)]_2$, R. Wilberger, H. Piotrowski, P. Mayer and I.-P. Lorenz, *Inorg. Chem. Commun.* **2002**, *5*, 897–902.
130. Naphtholactam als Komplexligand, M. Limmert, I.-P. Lorenz, J. Neubauer, A. Schulz und H. Piotrowski, *Z. Anorg. Allg. Chem.* **2003**, *629*, 223–231.

131. Synthesis of the dimeric, N substituted μ_2 -O aminoalkoxy complexes of rhenium $[(CO)_3Re(\mu_2-O^{\cap}NHR)]_2$, R. Wilberger, H. Piotrowski, P. Mayer, M. Vogt and I.-P. Lorenz, *Inorg. Chem. Commun.* **2003**, 6, 845–851.
132. Manganiophosphonium-Salze: Synthese, Eigenschaften und Struktur von Komplexen des Typs $[CpMn(CO)(NO)PPh_2H]BF_4$ und $[Cp'Mn(CO)(NO)PPh_2R]X$ (R = H, CH₃, [Cp'Mn(CO)(NO)]; X = BF₄, PF₆, S. Rudolph, J. Geicke, I.-P. Lorenz, T. Seifert, K. Polborn und M. Hörner, *Z. Naturforsch.* **2003**, 58b, 553–562.
133. Synthetic studies and structural aspects of some novel metallacyclic compounds of titanium(IV) incorporating nitrogen, oxygen and sulfur atoms: 1. Reactions of cis-dialkoxy-bis(acetylacetonato)titanium(IV) with alkoxyalkanols and crystal structure of a new modification of di- μ -oxo-bis(diacylacetone)titanium(IV)), M. Pathak, R. Bohra and R.C. Mehrotra, I.-P. Lorenz, H. Piotrowski, *Transition Metal Chemistry* **2003**, 28, 187–192.
134. Heteroleptic complexes of zirconium acetylacetones: Better precursors for the preparation of zirconia. Structural characterization of $[(acac)_2Zr\{ONC(Mepy-2)\}_2$, M. Pathak, R. Bohra, R.C. Mehrotra, I.-P. Lorenz and H. Piotrowski, *Z. Anorg. Allg. Chem.* **2003**, 629, 2493–2498.
135. Neue neutrale und kationische Mono-Aziridin-Komplexe des Typs $[CpMn(CO)_2Az]$, $[CpCr(NO)_2Az]^+$ und $[(Ph_3P)(CO)_4ReAz]^+$ über CO-, Hydrid- und Chlorid-Abspaltungsreaktionen, R. Wilberger, H. Piotrowski, M. Warchhold und I.-P. Lorenz, *Z. Anorg. Allg. Chem.* **2003**, 629, 2485–2492.
136. A new dichroic, nitroso bridged complex of rhenium: μ_2 -(η^2 -NO)-4-nitroso-*N,N*-dimethylaniline-bis[μ -chloro-tricarbonylrhenium(I)], R. Wilberger, C. Krinnerger, H. Piotrowski, P. Mayer, I.-P. Lorenz, *Eur. J. Inorg. Chem.* **2004**, 2488–2492.
137. Neutrale Bis-Aziridin-Komplexe des Typs $[M(CO)_3-XAz_2]$ (M = Me, Re; X = Cl, Br; Az = N(H)C₂H₂Me₂), R. Wilberger, C. Krinnerger, H. Piotrowski, P. Mayer, O. Ossberger, I.-P. Lorenz, *Z. Anorg. Allg. Chem.* **2004**, 630, 1495–1500.
138. Synthetic studies and structural aspects of some novel metallacyclic compounds of germanium(IV) incorporating nitrogen, oxygen and sulphur: 1 Reactions of germanium(IV) isopropoxide with diols and crystal structure of a unique diolato bridged germanium(IV) compound $[Ge_2(O_2C_2H_4)_4]$, M. Pathak, R. Bohra, R. C. Mehrotra, I.-P. Lorenz, H. Pietrowski, *J. Chem. Res.* **2004**, 567.
139. Reactions of $K_2[Fe(CO)_3PPh_3]$: Reductive Sb–Sb coupling with Ph₂SbCl to form *trans*-[$Fe(CO)_3PPh_3(-Sb_2Ph_4)$] and salt metathesis with Me₃SbCl₂ top to yield *trans*-[$Fe(CO)_3PPh_3(SbMe_3)$], I.-P. Lorenz, S. Rudolph, H. Piotrowski, K. Polborn, *Eur. J. Inorg. Chem.* **2005**, 82–85.
140. Synthesis and molecular structures of the cationic gold(I)-aziridine complexes $[Ph_3PAuAz]O_3SCF_3$ (Qz = C₂H₄NH, CH₂CHMeNH, CH₂CMe₂NH, CH₂CHe₂NH, CH₂CHPhNH, C₂H₄NBz, C₂H₄NC₂H₄OH), I.-P. Lorenz, C. Krinnerger, R. Wilberger, R. Bohra, H. Piotrowski, M. Warchhold, H. Nöth, *J. Organomet. Chem.* **2005**, 690, 1986–1993.

141. On the Electronic Structure and Bonding of the Intriguing Mixed $\text{Fe}^{2+}/\text{Fe}^{3+}$ $[\{\text{Cp}(\text{CO})_2\text{Fe}\}^t\text{BuPO}_2\text{FeCl}_2]_2$ -Complex, I.-P. Lorenz, M. Maier, A. Schulz, *Heteroatom Chem.* **2005**, *16*, 398–405.
142. Dichroic, dinuclear $\mu_2(\eta^2\text{-NO})$ -nitroso-aniline-bridged complexes of rhenium of the type $[(\text{CO})_3\text{ReX}]_2\text{ONC}_6\text{H}_4\text{NR}_2$ (X = Cl, Br, I, R = Me, Et), C. Krinninger, C. Högg, J.C.G. Ruiz, P. Mayer, O. Burkucky, A. Zumbusch, I.-P. Lorenz, *Chemistry* **2005**, *11*, 7228–7236.
143. Reactivity of 1-Nitroso-2-naphthol towards the Carbonyl Complexes $\text{XRe}(\text{CO}_5)$ (X = Al, Br, I): Molecular Structures of *cis*- $[\eta^2\text{-}\{\text{N},\text{O-C}_1\text{O}_6\text{ON(O)}\}\text{Re}(\text{CO})_4]$ and *fac*- $[\eta^2\text{-}\{\text{N},\text{O-C}_1\text{O}_6\text{ON(H)}\}\text{Re}(\text{CO})_3\text{X}]$, C. Krinninger, S. Wirth, J. C. G. Ruiz, P. Klüfers, H. Nöth, I.-P. Lorenz, *Eur. J. Inorg. Chem.* **2005**, 4094–4098.
144. Absence of dichroism in dinuclear rhenium complexes with sterically hindered $\mu_2(\eta^2\text{-N,O})$ -nitrosobenzene ligands, C. Krinninger, S. Wirth, P. Klüfers, P. Mayer, I.-P. Lorenz, *Eur. J. Inorg. Chem.* **2006**, 2060–2066.
145. Synthesis and structural elucidation of some novel aluminium(III) complexes with Schiff bases. Crystal and molecular structure of $[\text{Al}\{\text{O}(\text{C}_6\text{H}_4)\text{CH}=\text{NC}_6\text{H}_5\}_2\text{HO}(\text{C}_6\text{H}_4)\text{CH}=\text{NC}_6\text{H}_5\}_2]\text{Br}$, I.-P. Lorenz, M. Maier, A. Schulz, *Polyhedron* **2006**, 654–662.
146. Photolytisch induzierte Reaktionen von $\text{CpCo}(\text{CO})_2$ ($\text{Cp} = \text{C}_5\text{H}_5$, $\text{C}_5\text{H}_4\text{Me}$, C_5Me_5 und C_5Ph_5) mit Thiiran zu dinuklearen 1,2 Ethandithiolato-S,S-Komplexen, S. Drobnik, C. Stoll, H. Nöth, K. Polborn, W. Hiller, I.-P. Lorenz, *Z. Naturforsch.* **2006**, 61b, 1–12.
147. Synthesis, Cytotoxic Effect and Structure-Activity Relationship of Pd(II) Complexes with Coumarin Derivatives, Budzisz, E., Malecka, C., Lorenz, I.-P., Mayer, P., Kwiecien, R., Paneth, P., Krajewska, U., Rozalski, M., *Inorg. Chem.* **2006**, *45*, 9688–9695.
148. Synthesis, structures and comparison of neutral complexes formed by 2-(2'-pyridyl)indole and d6 transition metals, B. Neumann, C. Krinninger, I.-P. Lorenz, *European Journal of Inorganic Chemistry* **2007**, 472–480.
149. Glycol modified *cis*-diisopropoxy-bis(*N*-phenylsalicylideneiminato) titanium(IV). Syntheses, characterization, X-ray structures of $\text{Ti}\{\text{O}(\text{H}_6\text{H}_4)\text{CH}=\text{NPh}\}_2(\text{OPr}^i)_2$, trinuclear titanoxane $[\{\text{Ti}(\text{O}(\text{C}_6\text{H}_4)\text{CH}=\text{NPh})_2\}\text{O}]_3$ and their low temperature transformation to pure grains of nanosized titania, N. Sharma, V. Sharma, R. Bohra, V. S. Raju, I.-P. Lorenz, C. Krinninger and P. Mayer, *Inorg. Chim. Acta* **2007**, *360*, 3002–3012.
150. Synthesis and structural characterization of bis(aziridine) cobalt(II), zink(II) and palladium(II) complexes, J. N. Roedel, R. Bobka, M. Pfister, M. Rieger, B. Neumann and I.-P. Lorenz, *Z. Naturforsch.* **2007**, *62b*, 1095–1101.
151. Metal-mediated ring opening reaction of two aziridines to give 1,2-aminoaziridinoethane ligands, C. Krinninger, J. C. Ruiz, H. Noeth und I.-P. Lorenz, *Z. Anorg. Allg. Chemie* **2007**, *633*, 835–840.

152. Synthesis, characterization and structure of bis- and tetrakis-aziridine nickel(II) and copper(II) complexes, J. N. Roedel, R. Bobka, B. Neumann, B. Weber, P. Mayer and I.-P. Lorenz, *Z. Anorg. Allg. Chemie* **2007**, 633, 1171–1177.
153. Neutral mono- and cationic bis-aziridine d⁶-metal (π -arene)M(az)Cl₂ and [(π -arene)M(az)₂Cl]Cl (π -arene/M = η^6 -C₆Me₆/Ru, η^5 -C₅Me₅/Rh, Ir), R. Bobka, J. N. Roedel, B. Neumann, C. Krininger, P. Mayer, S. Wunderlich, A. Penger und I.-P. Lorenz, *Z. Anorg. Allg. Chemie* **2007**, 633, 1985–1994.
154. Spectroscopic and structural characterization of cationic *N*-(2-aminoethyl)aziridine-*N,N* chelate complexes of the d⁶-metals Rh(III) and Ir(III), R. Bobka, J. N. Roedel, B. Neumann, T. Nigst and I.-P. Lorenz, *Polyhedron* **2008**, 27, 955–961.
155. 2H-Azirines as ligands: Structural characterization of the first neutral and cationic η^5 -C₅Me₅Rh(III)-2H-azirine complexes, J. N. Roedel, I.-P. Lorenz, *Z. Anorg. Allg. Chem.* **2008**, 634, 1485–1489.
156. 2H-Azirines as ligands: Synthesis and Structural Characterization of (2H-3-arene-azirine)Re^I(CO)₃Br Complexes, J. N. Roedel, X. Wurzenberger, I.-P. Lorenz, *Inorg. Chem. Commun.* **2008**, 11, 829–831.
157. Cytotoxic effect, differentiation, inhibition of growth and theoretical calculations of an *N,N'*-donor ligand and its platinum(II), palladium(II) and copper(II) complexes, M. Miernicka, A. Szulawska, M. Czyz, I.-P. Lorenz, P. Mayer, B. Karwowski, E. Budzisz, *J. Inorg. Biochem.* **2008**, 102, 157–165.
158. Synthesis, crystal structure, theoretical calculation and cytotoxic effect of new Pt(II), Pd(II) and Cu(II) complexes with pyridine-pyrazoles derivatives, E. Budzisz, I.-P. Lorenz, P. Mayer, P. Paneth, L. Szatkowski, U. Krajewska, M. Rozalski and M. Miernicka, *New J. Chem.* **2008**, 32, 2238–2244.
159. Reactivity of 2,2-dimethylaziridine towards d¹⁰ transition metal halides, J. N. Roedel, F. Langenecker, I.-P. Lorenz, *J. Coord. Chem.* **2009**, 62, 1731–1742.
160. Tautomeric forms study of 1*H*-(2'pyridyl)-3-methyl-5-hydroxypyrazole and 1*H*-(2'-pyridil)-3-phenyl-5-hydroxypyrazole. Synthesis, structure and cytotoxic activity of their complexes with palladium(II) ion, M. Ciolkowski, P. Paneth, I.-P. Lorenz, P. Mayer, M. Rozalski, U. Krajewska, E. Budzisz, *J. Enz. Inhib. Med. Chem.* **2009**, 24, 1257–1268.
161. Synthesis and X-ray structure of platinum(II), palladium(II) and copper(II) complexes with pyridine-pyrazole ligands: Influence of ligands' structure on cytotoxic activity, E. Budzisz, M. Miernicka, I.-P. Lorenz, P. Mayer, U. Krajewska, M. Rozalski, *Polyhedron* **2009**, 28, 637–645.
162. Synthesis, Crystal Structure and Spectroscopic Characterization of Mono- and Di-nuclear 5,5-Diethylbarbiturato Complexes of Chromium(0) and Rhenium(I), N. Haque, J. N. Roedel, I.-P. Lorenz, *Z. Anorg. Allg. Chemie* **2009**, 633, 496–502.

163. Novel octahedral Bis{2-(aziridinyl)ethanol- κ^2N,O } Complexes of Cobalt(II), S. Putzien, S. Wirth, J. N. Roedel, I.-P. Lorenz, *Z. Anorg. Allg. Chem.* **2009**, 635, 1100–1105.
164. Di- and Triazaphospholes as Ligands for Ruthenium(II) and Iridium(III) Complexes, B. Köhler, S. Kammerer, T. Mayer, B. Neumann, I.-P. Lorenz, *Z. Anorg. Allg. Chem.* **2009** 635, 1362–1366.
165. Rhodium(III) and iridium(III) complexes with 1,2-naphthoquinone-1-oximate as bidentate ligand: synthesis, structure and biological activity, S. Wirth, C. J. Rohbogner, M. Cieslak, J. Kazmierczak-Baranska, S. Donevski, B. Nawrot, I.-P. Lorenz, *J. Biol. Inorg. Chem.* submitted **2009**, in press.
166. Tautomerization of 2-nitroso-N-arylanilines by coordination as *N,N'*-chelate ligands to rhenium(I) complexes and the anticancer activity of newly synthesized oximine rhenium(I) complexes against human melanoma and leukemia cells *in vitro*, S. Wirth, A. U. Wallek, A. Zernickel, M. Sztriller-Sikorska, K. Lesiak, I.-P. Lorenz, and M. Czyz, *J. Inorg. Biochem.* submitted **2009**.
167. Synthesis, structures, and characterization of benzildiimine complexes of rhodium(III) and iridium(I), N. Haque, B. Neumann, J. N. Roedel, I.-P. Lorenz, *Inorg. Chim. Acta* submitted **2009**, in press.
168. Formation of the complexes $[Fe(H_2BDI)_3]X_2$ and $[CpCr(H_2BDI)_2]X_2$ ($X = OTf$) from the unusual reactions of benzil-bis(trimethylsilyl)diimine with $CpML_2Cl$ ($M = Fe$, $L = CO$; $M = Cr$, $L = NO$), N. Haque, B. Neumann, I.-P. Lorenz, *Z. Naturforsch.* submitted **2009**, in press.
169. Synthesis and structures of Palladium(II) and Platinum(II) complexes with functional hydrazones, B. Köhler, S. Kammerer, L. Kovacevic, B. Neumann, I.-P. Lorenz, *Organometallics* submitted **2009**, in press.
170. Platinum-Assisted Isomerization of 1-Methylenehydrazine into 1-Carbenehydrazine, B. Köhler, I.-P. Lorenz, Th. M. Klapötke, *J. Phys. Chem. A* submitted **2009**, in press.